

SHORT NOTE

Changes in the number of black-billed gulls (*Larus bulleri*) nesting on the Ruamahanga River, Wairarapa, between 1998 and 2012

John M. Hill, Peter J. Bevan & John D. C. Bradley

Department of Conservation, Private Bag, Twizel 7901, New Zealand

Black-billed gulls (*Larus bulleri*) nesting on the Ruamahanga River, Wairarapa, New Zealand, were monitored between 1998 and 2012. The number of nesting gulls increased from 100 in 1998 to 1,200 in 2012. This increase was due to a combination of factors, including an increase in the number of nesting sites and an increase in the number of gulls nesting at each site. The increase in the number of nesting sites was due to the construction of a new nesting site in 2005. The increase in the number of gulls nesting at each site was due to an increase in the number of gulls nesting at the existing nesting sites. The increase in the number of gulls nesting at the existing nesting sites was due to an increase in the number of gulls nesting at the existing nesting sites. The increase in the number of gulls nesting at the existing nesting sites was due to an increase in the number of gulls nesting at the existing nesting sites.

Black-billed gulls (*Larus bulleri*) nesting on the Ruamahanga River, Wairarapa, New Zealand, were monitored between 1998 and 2012. The number of nesting gulls increased from 100 in 1998 to 1,200 in 2012. This increase was due to a combination of factors, including an increase in the number of nesting sites and an increase in the number of gulls nesting at each site. The increase in the number of nesting sites was due to the construction of a new nesting site in 2005. The increase in the number of gulls nesting at each site was due to an increase in the number of gulls nesting at the existing nesting sites. The increase in the number of gulls nesting at the existing nesting sites was due to an increase in the number of gulls nesting at the existing nesting sites.

Received 17 July 2015; accepted 19 January 2016
² arebergen@doc.govt.nz

Fig. 3. S, -Yd@Y-S a, -l
a, qdS .«" |lO«" ²C |«-l@
±S , 1" ±-l l±, qd/4
³²².². dYµ %Yµ±



©µ« Yµ±S ±a qd± fīl-° d"S
.« * dµd³ d 323 , d.±² ²C Yd@Y-S a, -l
.² Y qd, ±S YdS qdS, -l d±S
- 1" ±-l fīl ±². @±² ° «µ" «" a, -l ° 21"
², .lS " «" Yµ±S ±a qd± Y, -l ²C ±µl .²
±². « dS, µ±a qd, µl ± 1" Yµ qd/4 d.². d
²C Yd@Y-S a, -l ° µl ±, ±S ²± «" dle µ
l«² µl² «" Z d * dµd³ d | @-z | µ, µ pers.
comm . .« d. ± d Y. d l², 3-² d, -l « dS °
fī ±-l ¼Z d d±S -l 3² qdY. « d. «" a, -l l², ±S
²± Z d * dµd³ d ± ° -S 1" Yµ qd/4 ° µ" «"
% d d d a a % ± µ Yd@Y-S a, -l .l S ¼ l ± a
l²², µYd±S S YdS l ° ², S 3 µ 1" -Z d * dµd³ d
-l. «" ±² ± Yµ ± S ± a « Y- d @ µ. «" * dµd³ d Yd@
Y-S a, -l | «" Y l 3 d ± d±S . ± 2 | d | « d±S
l²², µYd±S . «" a, -l², .lS " «" Yµ ± S ± a |²² ± ¼
° ², S Y 3 µ ± ± ± a d fī ±-l ¼Z d
° d@Y-S a, -l dµ±" d fī ±-l ¼Z d ± d"
. a, -l d±S, 3. ² YdS l | d± Y. " ± µ ±
l³ µ ± a fī «" ¼µ. µ ± S ± «" ° S S ° ²C .
° ± µ ± ± ¼ d ° d | 2 ° YdS l dµ± S
d fī ±-l ¼Z d ²C «-l qd/4 S @µ d l², 3-
²C ° d, ±-l ² S ° d «" µ d±S l²² ° dS " «" °
- d1" d d ± µ, µ ± a ± , a, -l ¼ -S 1" Yµ qd/4
° ² l YdS l « d1" ° 3 µ d³ qd. 2 21" µ ± µ d Z d
* dµd³ d

«" dY- ¼ ²C Yd@Y-S a, -l .² ± l
l | l l ° ¼² ± «" % d d d a a % ± µ l a a l l
.« d l - dY- « Y- d µ l ± l ± d °, | « d. " µ S
µ l µ l ° / µ d µ " ± a. ² ± % a. ± d ° 2, ± l
- l d µ ²C . «" Yd@Y-S a, -l ²² ± ¼ d±S . «"
1 d " ²C * dµd³ d µ l µ @ µ YdS l " l ¼ d ¼
Yd±S d±S Yd@Qd ± S S ² µ l ± ±, d 22 S
l² ± µ ° 2 µ µ ° 21" qd ° 22 S ¼ 1" a. d. ± ± qd°
µ l µ l S l. «, l | µ d ± a d±S ° d ±. d ± ± a . «" 23 ±
« Y- d µ l µ YdS l µ " µ, ± °, S 2 ½ µ l | d± Y. l S
.² © µ « µ ± a d ± | µ l µ @ µ . «" Y ± " . ²C YdS l
d±S .². «" µ ° - S - C & µ d. «" qd ± a " | « d ± ± l d µ

²C - 1" d, .² YdS l d µ d µ l µ l S l z, -
| « d ± ± S µ l µ l ° « - d Yµ ± S ± a qd ± S l. « d
3 µ 1 - S " qd " 3 l. . ± ± d d ± l 3 µ S d µ l d±S qd
l µ C Qd ° ° ² qd 22 S ± a qd | µ | d. ² Yd@Y-S
a, -l d±S .². «" µ l µ µ YdS l

ACKNOWLEDGEMENTS

" «" d±S * dµd³ d µ l µ l µ l ° µ l ± a d S
Y ¼ µ l ° ± S d±S l 332 µ S d±S © ± S S Y ¼ / µ d µ
" ± a. ² ± % a. ± d ° 2, ± l - l 1 ²² S # l. . ± ± l . ± ±
± «" z d µ ± ± " | µ | 2 a ± ¼ a . «" d 32 µ d | ²C
* dµd³ d µ l µ l Yµ ± S ± a « Y- d @ µ ° d S µ l d±S Yd@
Y-S a, -l & l - d . d±S .² z d µ ± a ° 2 µ «" d l S
° 22 ± µ S z d µ l µ d ± d / ° % d±S z qd ° " ± a d
d 1 ° @ µ . «" µ l 332 µ .² . «" 3 l . . | « d ± d qd .²
µ l - l µ l . « d 3 µ 1 - S S |² ± l µ | - l | ² ° " ± l ² ± d ±
" d µ µ S µ d ° 2 C - l ° d³ d µ

LITERATURE CITED

- µ d ° L % fī d±S Yµ ± S ± a ²C Yd@Y-S a, -l ±
l . «" µ l fī d ° qd ° d ¼ d±S ±² µ «" µ * dµd³ d
Notornis 27
fī - a ± l # L. d - l qd & l l 1 " S l Handbook of Aus-
tralian, New Zealand & Antarctic birds. Volume 3, Snipe
to pigeons. z ° Y, µ " » © µ S (± l | µ l - ¼ µ l µ l
% Y µ ± ± µ ± µ YdS l, ° Y µ l ² ± µ ° d d d a a
* d ± a d ° * d qd ± ± d±S l d, «" µ l - d, µ l * d d
d d³ d ± l³ µ ± a 1² µ l ° ± S @ µ / µ d µ µ * -
± a. ² ± % a. ± d ° 2, ± l - 3
% Y µ ± ± µ ± µ l ² ± * dµd³ d % ± µ l d±S l² d l
± Yµ ± S ± a qd ± l² µ l ° ± S @ µ / µ d µ µ
* ± a. ² ± µ l a ± d ° 2, ± l - , ±³, Y l «" S µ 32 µ
3
& S 23 « % fī µ d@Y-S a, -l ±² µ «" µ * dµd³ d
µ d³ d Notornis 25

Keywords Yd@Y-S a, -l *Larus bulleri* 323, d.±²
| « d ± a Yµ ± S ± a |²² ± ¼ l